

КОГНИТИВНАЯ НАУКА В МОСКВЕ
НОВЫЕ ИССЛЕДОВАНИЯ



**МАТЕРИАЛЫ
КОНФЕРЕНЦИИ
2017**

ПОД РЕД. Е.В. ПЕЧЕНКОВОЙ, М.В. ФАЛИКМАН

УДК 159.9

ББК 81.002

К57

К57 Коллективный

Когнитивная наука в Москве: новые исследования. Материалы конференции 15 июня 2017 г.

Под ред. Е.В. Печенковой, М.В. Фаликман. – М.: ООО «Буки Веди», ИППИП. 2017 г. – 596 стр.

Электронная версия

ISBN 978-5-4465-1509-7

УДК 159.9

ББК 81.002

ISBN 978-5-4465-1509-7

© Авторы статей, 2017

TEST-RETEST RELIABILITY OF nTMS MULTI-MUSCLE MOTOR MAPPING

Nazarova M.A.* (1), Novikov P.A. (1, 2), Gusarovas I.V. (2),
Blagoveshchensky E.D. (1, 3), Nikulin V.V. (1, 4)
chantante@gmail.com

1 – Centre for Cognition and Decision Making, National Research University
Higher School of Economics, Moscow;

2 – Bauman Moscow State Technical University, Moscow;

3 – Laboratory of Neuroscience and Molecular Pharmacology, Institute
of Translational Biomedicine, St. Petersburg State University, St. Petersburg;

4 – Department of Neurology, Max Planck Institute for Human Cognitive and
Brain Sciences, Leipzig, Germany

Abstract. Despite being a routine technique for presurgical assessment, transcranial magnetic stimulation (TMS) mapping is underused for the probing of neuroplastic brain changes. We investigated the test-retest reproducibility of TMS cortical maps of several hand muscles using both standard and alternative parameters of the cortical representation in healthy volunteers. Pilot study results for four healthy right-handed male participants (19–33 y.o.) are presented. Two TMS mapping sessions with stimulation of the left motor cortex were performed within 5 to 10 days (Day1 and Day2). Day2 points repeated an exact order of the Day1 session. For quantitative comparison of the 3D profile similarities, earth mover's distance metrics was used. Analysis of nTMS maps was performed using the custom-made software TMSmap (<http://tmsmap.ru>). The between-days difference in the area of cortical representation for the four analyzed participants was 14.5–30.4% for one repetition and 3.9–11.2% for five repetitions of each cortical point. Considering the 3D profiles of cortical representation, higher similarity was shown for the same muscles' representations and their overlaps compared to the representations of the different muscles. The study is ongoing, and further analyzed results will be presented as they become available.

Keywords: motor cortex, transcranial magnetic stimulation, functional brain mapping

The study has been partly funded by the Russian Academic Excellence Project '5-100' and RFBR grant № 16-04-01883.

Background

Navigated-TMS (nTMS) motor mapping has become a routine technique for presurgical cortical assessment (Picht et al., 2016; Tarapore et al., 2016). However, nTMS mapping to investigate cortical neuroplastic changes during rehabilitation, training or for any other longitudinal research is still limited both in clinical and basic research. This is largely due to the lack of any generally accepted methodology of nTMS mapping (Lüdemann-Podubecká, Nowak, 2016). Thus,

there is no agreement on how many stimulation points and how many repetitions for the each point are necessary and which parameters of TMS maps besides the area are useful for reliable mapping. Moreover, it is not clear to what extent motor maps are reproducible even in healthy participants, which seriously affects the interpretation of dynamic changes with an nTMS mapping approach. In this work, our aim was to investigate the test-retest reproducibility of the nTMS cortical maps of the upper-limb muscles in healthy volunteers using both standard and several new parameters such as the overlap between the cortical representation of different muscles (Nazarova, 2015) and 3D profiles of the cortical motor maps considering the motor evoked amplitudes (MEP) in each point (Raffin et al., 2015).

Method

Here we present the results of the pilot study performed on four healthy right-handed male volunteers (19–33 y.o.). Two TMS mapping sessions with stimulation of the left motor cortex were performed within 5 to 10 days (Day1 and Day2), with an investigation of the cortical representations of three upper limb muscles: the abductor pollicis brevis (APB), abductor digiti minimi (ADM), and extensor digitorum communis (EDC). Sessions consisted of 60–90 points stimulated in a pseudo-random order five times (see Fig. 1). Day 2 points were stimulated in the exact same order as on Day 1. Analysis was performed using the custom-made software TMSmap (<http://tmsmap.ru/>) (see Fig. 2). For quantitative comparison of the 3D profile similarities, we used earth mover's distance metrics (EMD), the so called Wasserstein metric.

Results

Below we present the initial results. The four analyzed participants demonstrated higher reproducibility of the maps in the case of five repetitions of each point compared to only one stimulation of each point. The difference in the area of cortical representation for Day1 and Day2 was 14.5–30.4% for one stimulation and only 3.9–11.2% for five repetitions, showing that the commonly used




Figure 1. nTMS mapping procedure: 5 sub-sessions in the first mapping day (Day1) constitute the whole session. Five coordinates (x, y, z) and MEP-amplitudes in each target spot related to one node of the grid were averaged (black arrow)




Figure 2. Muscle cortical representations stability from Day1 to Day2 (Subj. 2).

approach for presurgical mapping (one stimulation per site) may be less applicable for longitudinal studies. Considering the 3D profiles of the cortical representations, higher similarity was shown for the same muscles' representations and their overlaps compared to the representations of different muscles based on the EMD metric, reflecting that the 3D profiles of the cortical representation considering MEP amplitudes in each point may be functionally meaningful and not just noise as it was regarded before. The obtained results provide further evidence for the usefulness of the more elaborate metrics such as 3D profiles and the overlaps between muscle cortical representations when estimating the stability of motor TMS maps. More detailed results will be presented in the future.

References

- Lüdemann-Podubecká J., Nowak D.A. Mapping cortical hand motor representation using TMS: A method to assess brain plasticity and a surrogate marker for recovery of function after stroke? // *Neuroscience and Biobehavioral Reviews*. 2016. Vol.69. P. 239–51. [doi:10.1016/j.neubiorev.2016.07.006](https://doi.org/10.1016/j.neubiorev.2016.07.006)
- Nazarova M.A. Multimodal assessment of the motor system reorganization after ischemic hemispheric stroke: MRI-TMS study. PhD dissertation. Research Center of Neurology of Russian Academy of Med. Sciences, Moscow, 2015.

Picht T., Frey D., Thieme S., Kliesch S., Vajkoczy P. Presurgical navigated TMS motor cortex mapping improves outcome in glioblastoma surgery: A controlled observational study // Journal of Neuro-Oncology. 2016. Vol. 126. No. 3. P. 535 – 43. doi:10.1007/s11060-015-1993-9

Raffin E., Pellegrino G., Lazzaro V.D., Thielscher A., Siebner H.R. Bringing transcranial mapping into shape: Sulcus-aligned mapping captures motor somatotopy in human primary motor hand area // NeuroImage. 2015. Vol. 120. P. 164 – 75. doi:10.1016/j.neuroimage.2015.07.024

Tarapore P.E., Picht T., Bulubas L., Shin Y., Kulchytska N., Meyer B., Berger M.S., Nagarajan S.S., Krieg S.M. Safety and tolerability of navigated TMS for preoperative mapping in neurosurgical patients // Clinical Neurophysiology. 2016. Vol. 127. No. 3. P. 1895 – 1900. doi:10.1016/j.clinph.2015.11.042

Стабильность двигательных корковых репрезентаций при навигационной ТМС моторной коры

М.А. Назарова* (1), П.А. Новиков* (1, 2), И.В. Гусаровас (2),
Е.Д. Благовещенский (1, 3), В.В. Никулин (1, 4)
chantante@gmail.com

1 – Центр нейроэкономики и когнитивных исследований, НИУ «Высшая школа экономики», Москва;

2 – МГТУ им. Н.Э. Баумана, Москва;

3 – Лаборатория нейробиологии и молекулярной фармакологии, Институт трансляционной биомедицины, Санкт-Петербургский государственный университет, Санкт-Петербург;

4 – Department of Neurology Max Planck Institute for Human Cognitive and Brain Sciences, Лейпциг, Германия

Аннотация. Предхирургическое картирование моторных зон коры с помощью навигационной транскраниальной магнитной стимуляции (нТМС) является на настоящий момент достаточно распространенным клиническим подходом. В данной работе мы исследовали стабильность различных параметров нТМС корковых репрезентаций нескольких мышц руки, включая такие новые параметры как 3D профили ТМС карт и их наложение. Представлены результаты пилотного исследования четырех здоровых добровольцев (19–33 года). Исследование включало две повторные ТМС сессии левой двигательной коры, проведенные с разницей в 5–10 дней (День1 и День2). Анализ проводился с использованием разработанной программы TMSmap (<http://tmsmap.ru/>). Для количественной оценки 3D профилей использовалась метрика Вассерштейна (earth mover's distance). Предварительные результаты для четырех добровольцев показали большую стабильность стандартных показателей карт при пяти повторениях стимуляции каждой точки по сравнению с единичной стимуляцией. Так, разница площадей корковых репрезентаций для сессий первого и второго дня составила 14.5–30.4% в случае единичного повторения и 3.9–11.2% в случае пяти повторений. Относительно 3D профилей корковых репрезентаций большая схожесть карт День1 – День2 (меньшее значение EMD) было показано для репрезентаций одних и тех же мышц и их наложений по сравнению с корковыми картами различных мышц. Полученные результаты подчеркивают важность использования дополнительных параметров для оценки стабильности двигательных ТМС карт.

Ключевые слова: двигательная кора, навигационная транскраниальная магнитная стимуляция, функциональное картирование мозга